

Constituent Quarks: The Beginning of the End

G. Zweig, BNL
January 29, 2013

- QCD - developed in two phases:
 - Discovery of quarks
 - Specification of their interactions
- Arose from two very different traditions
 - Rutherford-Bohr
 - Einstein
- Discovery of radioactivity: Henri Becquerel (1896)



Figure 1: Becquerel's photographic plate fogged by exposure to radiation from uranium salts. The shadow of a metal Maltese Cross placed between the plate and the uranium salts is visible.

- Rutherford at Cambridge (1899): α and β

- Rutherford & Soddy at McGill (1903):
“disintegration theory” of radioactivity
radioactivity an atomic - not molecular - process

Nobel prize in Chemistry (1908)



Figure 2: Rutherford's group at Manchester University, 1910

PROCEEDINGS OF
THE ROYAL SOCIETY.

SECTION A.—MATHEMATICAL AND PHYSICAL SCIENCES.

On a Diffuse Reflection of the α -Particles.

By H. GEIGER, Ph.D., John Harling Fellow, and E. MARSDEN, Hatfield
Scholar, University of Manchester.

(Communicated by Prof. E. Rutherford, F.R.S. Received May 19,—Read
June 17, 1909.)

In the following experiments, however, conclusive evidence was found of the existence of a diffuse reflection of the α -particles. A small fraction of the α -particles falling upon a metal plate have their directions changed to such an extent that they emerge again at the side of incidence. To form an

We are indebted to Prof. Rutherford for his kind interest and advice throughout this research.

- Impossible
- Marsden (1914): Nuclei contain protons
- Bohr (1914-1916): Rutherford plus Planck
- Heisenberg (1925, 1943, & 1944): Work only with observables.

Lone Ranger Atom Bomb Ring Spinthariscopes (1947 - early 1950s)

This ring spinthariscopes was known as the Lone Ranger Atom Bomb Ring and advertised as a "seething scientific creation." The Lone Ranger was more closely associated with silver bullets than atomic bombs but that's what it was called. When the red base (which served as a "secret message compartment") was taken off, and after a suitable period of time for dark adaptation, you could look through a small plastic lens at scintillations caused by polonium alpha particles striking a zinc sulfide screen.



Distributed by Kix Cereals (15 cents plus a boxtop), the instructions stated: "You'll see brilliant flashes of light in the inky darkness inside the atom chamber. These frenzied vivid flashes are caused by the released energy of atoms. PERFECTLY SAFE - We guarantee you can wear the KIX Atomic "Bomb" Ring with complete safety. The atomic materials inside the ring are harmless."

The following advertisement was appearing in newspapers in early 1947.

Advertisement
Advertisement

SEE GENUINE ATOMS SPLIT TO SMITHEREENS!

INSIDE THIS KIX ATOMIC "BOMB" RING!

HAS SECRET MESSAGE COMPARTMENT

Only 15¢ Plus Boxtop

Kix

How Tommy Thwarted the Enemy Agents

with his KIX Atomic "Bomb" Ring

IN THE TIME KIX RING NOVEL, THE EFFECTS OF ATOMIC ENERGY IN ACTION! IT'S AS FANTASY AS OUR NEW SECRET FORMULA!

SEE, THE ENEMY IS SWARMING AND, BUT TOMMY KNOWS SECRET! HE'S USING THE TOP-SECRET ATOMIC FORMULA IN THIS KIX RING TO REVEAL THE TRUTH!

BUT-HAT CHIMP! HEADQUARTERS! I LEARNED THAT A KIX RING WAS CAUSING THE ENEMY TO REVEAL THE TRUTH!

THEY WERE TOMMY'S TOP-SECRET ATOMIC FORMULA AND FORMULA WILL HELP REVEAL THE TRUTH!

SEE, GENERAL! THE ENEMY'S SECRET MESSAGE WAS HIDDEN IN THE KIX RING! IT WAS REALLY FOOLING THE ENEMY!

HERE'S HOW TOMMY WAS ABLE TO THWART THE ENEMY AGENTS --

FORMULA WAS HIDDEN IN SECRET MESSAGE COMPARTMENT INSIDE TOP OF KIX ATOMIC "BOMB" RING

WITHIN ATOM CHAMBER IN THUMBHEAD

CONCEALED OBSERVATION LENS

It's a SEETHING SCIENTIFIC SENSATION!

Actual Atoms—right in smoldering inside this ring! And you can see brilliant atomic effects! Take ring into dark room and wait until your eyes are accustomed to darkness. Slide Tail-Fin off—look in Observation Lens—and you'll see frenzied flashes of light—caused by released energy of atoms splitting like crazy. Secret Message Compartment hidden in Tail Fin... Bombardier's insignia embossed on top of ring. Slide silver shield lighting-blast explosions. Aluminum Warhead (bomb nose)... 4-pronged plastic Tail Fin. Ring can be adjusted to fit any finger. Atomic materials in Atom Chamber are harmless. Now is Got Ring. Get a line of that winning corn-daddy breakfast cereal KIX. Bend top of KIX box together with only life and Order Blank at right.

HURRY—MAIL THIS ORDER BLANK TODAY!

SEND TO: KIX, Box 65, New York City, N. Y.

Here's my 15¢ plus KIX boxtop. Back to me at once my KIX Atomic "Bomb" Ring.

NAME.....

ADDRESS.....

CITY.....

State.....

First names and address please. Use your number (if any).

NOTE: If you prefer the new "KIX Atomic Bomb" ring package with silver black on top, use that order blank instead of this.

Offer good in U.S.A. only, while supplies last.

General Mills

THE PHYSICAL REVIEW

A journal of experimental and theoretical physics established by E. L. Nichols in 1893

SECOND SERIES, VOL. 76, No. 12

DECEMBER 15, 1949

Are Mesons Elementary Particles?

E. FERMI AND C. N. YANG*

Institute for Nuclear Studies, University of Chicago, Chicago, Illinois

(Received August 24, 1949)

The hypothesis that π -mesons may be composite particles formed by the association of a nucleon with an anti-nucleon is discussed. From an extremely crude discussion of the model it appears that such a meson would have in most respects properties similar to those of the meson of the Yukawa theory.

I. INTRODUCTION

IN recent years several new particles have been discovered which are currently assumed to be "elementary," that is, essentially, structureless. The probability that all such particles should be really elementary becomes less and less as their number increases.

It is by no means certain that nucleons, mesons, electrons, neutrinos are all elementary particles and it could be that at least some of the failures of the present theories may be due to disregarding the possibility that some of them may have a complex structure. Unfortunately, we have no clue to decide whether this is true, much less to find out what particles are simple and what particles are complex. In what follows we will try to work out in some detail a special example more as an illustration of a possible program of the theory of particles, than in the hope that what we suggest may actually correspond to reality.

We propose to discuss the hypothesis that the π -meson may not be elementary, but may be a composite particle formed by the association of a nucleon and an anti-nucleon. The first assumption will be, therefore, that both an anti-proton and an anti-neutron exist, having the same relationship to the proton and the neutron, as the electron to the positron. Although this is an assumption that goes beyond what is known experimentally, we do not view it as a very revolutionary one. We must assume, further, that between a nucleon and an anti-nucleon strong attractive forces exist, capable of binding the two particles together.

* Now at the Institute for Advanced Study, Princeton, New Jersey.

We assume that the π -meson is a pair of nucleon and anti-nucleon bound in this way. Since the mass of the π -meson is much smaller than twice the mass of a nucleon, it is necessary to assume that the binding energy is so great that its mass equivalent is equal to the difference between twice the mass of the nucleon and the mass of the meson.

According to this view the positive meson would be the association of a proton and an anti-neutron and the negative meson would be the association of an anti-proton and a neutron. As a model of a neutral meson one could take either a pair of a neutron and an anti-neutron, or of a proton and an anti-proton.

It would be difficult to set up a not too complicated scheme of forces between a nucleon and an anti-nucleon, without about equally strong forces between two ordinary nucleons. These last forces, however, would be quite different from the ordinary nuclear forces, because they would have much greater energy and much shorter range. The reason why no experimental indication of them has been observed for ordinary nucleons may be explained by the assumption that the forces could be attractive between a nucleon and an anti-nucleon and repulsive between two ordinary nucleons. If this is the case, no bound system of two ordinary nucleons would result out of this particular type of interaction. Because of the short range very little would be noticed of such forces even in scattering phenomena.

Ordinary nuclear forces from the point of view of this theory will be discussed below.

Unfortunately we have not succeeded in working out a satisfactory relativistically invariant theory of nucleons among which such attractive forces act. For this reason all the conclusion that will be presented will be

- M. Gell-Mann & E.P. Rosenbaum, “Elementary Particles,” *Scientific American*, July 1957, 72-86: 19 in number

M. Gell-Mann & A.H. Rosenfeld, “Hyperons and Heavy Mesons,” *Ann. Rev. Nucl. Sci*, 1957, 407-478:

Point particles

Spin 1/2 leptons	
Particle	Mass
e^-	1
μ^-	206.7
ν	0

Spin 1 photon	
Particle	Mass
γ	0

Extended particles (strongly interacting)

Spin 1/2 baryons		
Multiplet	Particle	Mass (m_e)
Ξ	Ξ^0	?
	Ξ^{-1}	2585
Σ	Σ^{-1}	2341
	Σ^+	2325
	Σ^0	2324
Λ	Λ	2182
N	n	1838.6
	p	1836.1

Spin 0 mesons		
Multiplet	Particle	Mass
π	π^+	273.2
	π^{-1}	273.2
	π^0	264.2
K	K^+	966.5
	K^-	966.5
	K_1^0	965
	K_2^0	965

– No resonances mentioned!





FIG. 3. A plot of $\pi^\pm + p$ total cross-sections as a function of energy, using in general

- Caltech:
 - My thesis: A test of time reversal symmetry
 $K^+ \rightarrow \pi^0 + \mu^+ + \nu$.
 - Mexico!
 - Murray?
- Every Thursday at 1:30 PM during 1962-63
- Theoretical physics
 - Axiomatic field theory
 - Theory related to belief (Chew, June 1961):
 “I believe the conventional association of fields with strongly interacting particles to be empty. ... field theory..., like an old soldier, is destined not to die but just fade away.”
 - Theory related to experiment:
 - * Particle classification (no dynamics)

* Dynamics - bootstrap (no classification)



The ρ bootstrap, F. Zachariasen, *Phys. Rev. Lett.* **7**, 112 (1961)

- Experimental physics

- More particles:

- * A 4th lepton (ν)

- * An 8th spin 0 meson (η)

- * Many more resonances: RMP March 1963, ρ , ω , K^* , \dots (26 meson resonances listed)

- One Thursday afternoon:

P.L. Connolly, et al., “Existence and Properties of the ϕ Meson”, *Phys. Rev. Lett.* **10**, 371 (1963):




FIG. 1. Dalitz plot for the reaction $K^- + p \rightarrow \Lambda + K + \bar{K}$. The effective-mass distribution for $K\bar{K}$ and for ΛK^+ are projected on the abscissa and ordinate (see reference 7).

$$\phi \rightarrow K \bar{K}$$




FIG. 4. The $M(\pi^+\pi^-\pi^0)$ distribution from the reaction $K^- + p \rightarrow \Lambda + \pi^+ + \pi^- + \pi^0$ after removing Y_1^* production events (see text).

$$\phi \not\rightarrow \rho + \pi$$

$$\begin{aligned}
\frac{\Gamma_{K\bar{K}}}{\Gamma_{\rho\pi}} &\sim \left(\frac{p_{K\bar{K}}}{p_{\rho\pi}} \right)^3, \\
&= 0.23 \text{ (expected),} \\
&\geq 35 \text{ (observed).}
\end{aligned}$$

“The observed rate [for $\phi \rightarrow \rho + \pi$] is lower than ... predicted values by one order of magnitude; however the above estimates are uncertain by at least this amount so that this discrepancy need not be disconcerting.”

— Feynman:


- Assume hadrons have constituents called aces:

$$[(p_0, n_0), \Lambda_0] \quad \& \quad [(\bar{p}_0, \bar{n}_0), \bar{\Lambda}_0]$$



Vector mesons as “deuces”, CERN report TH-401, January 1964

- A rule for *dynamics*:



Meson decay: a is an ace, \bar{a} an antiace.

– Implies $\phi \not\rightarrow \rho + \pi$

More specifically:

- Hadrons have *point* constituents
- Lepton \leftrightarrow ace correspondance
- Unknown number: represented by regular polygons of increasing size
- Origin of SU(3) symmetry
- Beyond SU(3) symmetry:
 - Restricted representations, quantum numbers:
 - * Baryons only in 1,8,10;
mesons only in 1,8, and 9
 - * $L = 0$ baryons: $(8, J^P = \frac{1}{2}^+)$ & $(10, \frac{3}{2}^+)$;
 $L = 0$ mesons: $J^{PC} = 0^{-+}$ and 1^{--} .
 - * $0^{--}; 0^{+-}, 1^{-+} \dots$ forbidden for any L
 - Symmetries broken by ace mass splittings
 - * SU(3): $m(p_0) = m(n_0) < m(\Lambda_0)$
 - * SU(2): $m(p_0) < m(n_0)$


$$\begin{aligned}
p &= \frac{1}{\sqrt{2}} \left(\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \blacktriangle \quad \bullet \end{array} - \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \blacktriangle \end{array} \right) & n &= \frac{1}{\sqrt{2}} \left(\begin{array}{c} \blacktriangle \\ \diagup \quad \diagdown \\ \blacktriangle \quad \bullet \end{array} - \begin{array}{c} \blacktriangle \\ \diagup \quad \diagdown \\ \bullet \quad \blacktriangle \end{array} \right) \\
\Lambda &= \frac{1}{\sqrt{12}} \left(\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \blacksquare \quad \blacktriangle \end{array} - \begin{array}{c} \blacktriangle \\ \diagup \quad \diagdown \\ \blacksquare \quad \bullet \end{array} + \begin{array}{c} \blacktriangle \\ \diagup \quad \diagdown \\ \bullet \quad \blacksquare \end{array} - \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \blacktriangle \quad \blacksquare \end{array} + 2 \begin{array}{c} \blacksquare \\ \diagup \quad \diagdown \\ \bullet \quad \blacktriangle \end{array} - 2 \begin{array}{c} \blacksquare \\ \diagup \quad \diagdown \\ \blacktriangle \quad \bullet \end{array} \right) \\
\Sigma^0 &= \frac{1}{2} \left(\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \blacksquare \quad \blacktriangle \end{array} + \begin{array}{c} \blacktriangle \\ \diagup \quad \diagdown \\ \blacksquare \quad \bullet \end{array} - \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \blacktriangle \quad \blacksquare \end{array} - \begin{array}{c} \blacktriangle \\ \diagup \quad \diagdown \\ \bullet \quad \blacksquare \end{array} \right) \\
\Sigma^+ &= \frac{1}{\sqrt{2}} \left(\begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \blacksquare \end{array} - \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \blacksquare \quad \bullet \end{array} \right) & \Sigma^- &= \frac{1}{\sqrt{2}} \left(\begin{array}{c} \blacktriangle \\ \diagup \quad \diagdown \\ \blacksquare \quad \blacktriangle \end{array} - \begin{array}{c} \blacktriangle \\ \diagup \quad \diagdown \\ \blacktriangle \quad \blacksquare \end{array} \right) \\
\Xi^0 &= \frac{1}{\sqrt{2}} \left(\begin{array}{c} \blacksquare \\ \diagup \quad \diagdown \\ \bullet \quad \blacksquare \end{array} - \begin{array}{c} \blacksquare \\ \diagup \quad \diagdown \\ \blacksquare \quad \bullet \end{array} \right) & \Xi^- &= \frac{1}{\sqrt{2}} \left(\begin{array}{c} \blacksquare \\ \diagup \quad \diagdown \\ \blacksquare \quad \blacktriangle \end{array} - \begin{array}{c} \blacksquare \\ \diagup \quad \diagdown \\ \blacktriangle \quad \blacksquare \end{array} \right)
\end{aligned}$$

FIG. 3

Hadrons as “treys”, CERN report TH-412, February 1964

– *Rule #1: Aces, not hadrons, interact.*

* Strong interactions: “Zweig’s rule” (what’s allowed!)



Graphical representation of the meson-baryon coupling.

The “little loop” encloses antisymmetrized aces.

The subscript “0” on aces is suppressed.

* Electromagnetic and weak interactions:

$$a \rightarrow a + \gamma$$

$$a \rightarrow a' + e^- + \nu$$

– *Rule #2: Mass = Constituent masses + pairwise energies of interaction,*
 $|\Delta m| > |\Delta E|.$

* Mesons:

$$m(D_{\alpha}^{\beta}) = m(\alpha) + m(\beta) - E_{\alpha}^{\beta}$$

* Baryons:

$$m(T_{\alpha\beta\gamma}^8) = m(\alpha) + m(\beta) + m(\gamma) - E_{\alpha\beta.}^8 - E_{\alpha.\gamma}^8 - E_{.\beta\gamma}^8$$

No potential function assumed.

Not the naive quark model!

- Would you have believed?

- SU(3) symmetry breaking:

$$m(a_1) = m(a_2) < m(a_3)$$

Hierarchy of baryon relations, e.g., $\frac{1}{2}^+$ octet:

$$* E_{\alpha\beta.}^8 \approx E_{\gamma\delta.}^8 \dots$$

$$\cdot \begin{array}{cccc} m(N) < m(\Lambda) & \approx & m(\Sigma) < m(\Xi) \\ 939 & 1115 & 1193 & 1316 \end{array}$$

$$\cdot \begin{array}{ccc} \frac{1}{2}[m(\Xi) + m(N)] & \approx & \frac{1}{2}[m(\Lambda) + m(\Sigma)] \\ 1127 & & 1154 \end{array}$$

$$* \frac{1}{2}(E_{33.}^8 + E_{\alpha\beta.}^8) \approx E_{3\alpha.}^8 \approx E_{3\beta.}^8, \alpha, \beta = 1, 2$$

$$\cdot \begin{array}{ccc} \frac{1}{2}[m(\Xi) + m(N)] & \approx & \frac{1}{4}[3m(\Lambda) + m(\Sigma)] \\ 1127 & & 1134 \end{array}$$

(Gell-Mann Okubo mass formula)

Hierarchy of meson relations, e.g.,
 $J^{PC} = 1^{--}$ nonet:

$$* E_{\alpha}^{\beta} \approx E_{\gamma}^{\delta} \dots,$$

$$\cdot \begin{array}{cccc} m(\rho) & \approx & m(\omega) & < & m(K^*) & < & m(\phi) \\ 750 & & 784 & & 888 & & 1018 \end{array}$$

$$* \frac{1}{2}(E_3^3 + E_{\alpha}^{\beta}) \approx E_3^{\beta} \approx E_{\alpha}^3, \quad \alpha, \beta = 1, 2$$

$$\cdot \begin{array}{ccc} m^2(\phi) & \approx & 2m^2(K^*) - m^2(\rho) \\ (1018)^2 & & (1007)^2 \end{array}$$

– SU(2) breaking:

$$m(a_1) < m(a_2)$$

Hierarchy of baryon relations, e.g., $\frac{1}{2}^+$ octet:

$$* E_{\alpha\beta}^8 \approx E_{\alpha'\beta'}^8 \dots$$

$$\cdot \begin{array}{cc} m(p) < m(n) \\ 938 & 939 \end{array}$$

$$\cdot \begin{array}{ccc} m(\Sigma^+) < m(\Sigma^0) < m(\Sigma^-) \\ 1190 & 1193 & 1196 \end{array}$$

$$\cdot \begin{array}{cc} m(\Xi^0) < m(\Xi^-) \\ 1315 & 1321 \end{array}$$

$$* \frac{1}{2}(E_{11}^8 + E_{22}^8) \approx E_{12}^8, \dots$$

$$\cdot \begin{array}{ccc} \frac{1}{2}[m(\Sigma^+) + m(\Sigma^-)] \approx m(\Sigma^0) \\ 1193.4 \pm 0.3 & & 1193.2 \pm 0.7 \end{array}$$

* No interaction energy assumptions

$m(n) - m(p) + m(\Xi^-) - m(\Xi^0) = m(\Sigma^-) - m(\Sigma^+)$	
7.3 ± 1.3	8.3 ± 0.5
$\{8.14 \pm 0.21\}$	$\{8.08 \pm 0.08\}$
$\{\Delta = 0.06 \pm 0.22\}$	

- Additional relations for 0^{-+} meson octet and baryon $\frac{3}{2}^{+}$ decuplet
- Cross multiplet relations ($\Delta m > \Delta E$)
 - * $m(\Xi^{*}) - m(\Sigma^{*}) \approx m(\Xi) - m(\Sigma)$

145
130
 - * $m^2(K^{*}) - m^2(\rho) \approx m^2(K) - m^2(\pi)$

0.22 GeV²
0.22 GeV²
 - * $m(N) < m(\Lambda) \Rightarrow m(\rho) < m(K^{*})$
- Other baryons & mesons with $L \neq 0$ ($\vec{L} \cdot \vec{S}$)
- Overfitting?
 - [QCD \Rightarrow ?]
- Not as easy as it looks
 - * 26 \Rightarrow 7, exotics
 - * An earlier Thursday afternoon

- What did people think?
 - GZ: Duck test
 - MGM:
 - * “Concrete quark model”
 - * Five years after the deep inelastic scattering experiments at SLAC (partons) “Quarks,” *Acta Physica Austriaca, Suppl. IX*, 733-761 (1972)

“In these lectures I want to speak about at least two interpretations of the concept of quarks for hadrons and, the possible relations between them.

First I want to talk about quarks as ‘constituent quarks’. These were used especially by G. Zweig (1964) who referred to them as aces. ...

The whole idea is that hadrons act as if they are made up of quarks, but the quarks do not have to be real. If we use the quark statistics described above, we see that it would be hard to make the quarks real ...

There is a second use of quarks, as so-called ‘current quarks’ which is quite different from their use as constituent quarks ...

If quarks are only fictitious there are certain defects and virtues. The main defect would be that we never experimentally discover real ones and thus will never have a quarkonics industry. The virtue is that then there are no basic constituents for hadrons — hadrons act as if they were made up of quarks but no quarks exist - and, therefore, there is no reason for a distinction between the quark and bootstrap picture: they can be just two different descriptions of the same system, like wave mechanics and matrix mechanics.”

— RPF:

- * Current quarks?
- * Constituent quarks, big problem!
- * The correct theory should not allow one to say which particles are elementary.
- * “Have I missed anything Zweig?”

Problems with acceptance:

- Aces impossible according to the spin-statistics theorem.
 - * Rutherford's atom & Bohr's orbits
 - * Wegener's continental drift
- Aces impossible according to current dogma:
 1. Aces incompatible with nuclear democracy,
 2. The rules for aces incompatible with Heisenberg's observability requirement.
 - * Copernicus's view of the solar system

Acceptance criteria for ace constituents:

Bayes Theorem:

$$P(A) \ll 1, \quad P(E|A) \approx 1.$$

$$P(A|E) \approx \frac{1}{1+\lambda},$$

$$\lambda \equiv \frac{P(E|\bar{A})}{P(A)}.$$

Acceptance when $P(E|\bar{A}) \ll P(A)$.

- Einstein tradition: $P(E|\bar{A}) \gg P(A)$:
- Rutherford-Bohr tradition: $P(E|\bar{A}) \ll P(A)$

When did acceptance come?

- Pauling
- Dalitz
- Feynman
- Deep inelastic scattering
- ψ/J

Invention or discovery?

- Current quarks invented (Einstein tradition)
- Constituent quarks (aces) discovered (Rutherford-Bohr tradition)

There are, however, many unanswered questions. Are aces particles ? If so, what are their interactions ? Do aces bind to form only deuces and treys ? What is the particle (or particles) that is responsible for binding the aces ? Why must one work with masses for the baryons and mass squares for the mesons ? And more generally, why does so simple a model yield such a good approximation to nature ?

there is also the outside chance that the model is a closer approximation to nature than we may think, and that fractionally charged aces abound within us.

Conclusion, CERN report TH-412, February 1964